

D-Stability and Real and Complex Quadratic Forms

Charles R. Johnson*

National Bureau of Standards, Washington, D.C. 20234[†]

Recommended by David H. Carlson

ABSTRACT

The problem of characterizing the class of D -stable matrices has remained unsolved since its suggestion in a paper by Arrow and McManus [1] in 1958. In this note we present a necessary condition involving real quadratic forms which is not sufficient and a sufficient condition involving complex quadratic forms which is not necessary.

Denote by $M_n(F)$ the class of n by n matrices over the field F , where F may be either the real numbers R or the complex numbers C . Let D_n be the class of diagonal matrices with real positive diagonal entries. We denote the spectrum of $A \in M_n(F)$ by $\sigma(A)$. Then $A \in M_n(F)$ is said to be D -stable if $D \in D_n$ implies $\operatorname{Re}(\lambda) > 0$ for all $\lambda \in \sigma(DA)$. We shall denote the class of D -stable elements of $M_n(F)$ by $DL_n(F)$ and their Euclidean closure by $\overline{DL_n(F)}$.

We shall not review the known facts about D -stability [1, 6, 7, 8, 9] except to mention that the broadest known sufficient condition is as follows. If there is a $D \in D_n$ such that $DA + A^*D$ is positive definite, then $A \in DL_n(C)$. Several of the other sufficient conditions are known to be special cases of this one [4, 5, 7]. However, this condition is a special case of the sufficient condition given by Theorem 2 below.

We proceed to relate D -stability to quadratic forms. Theorem 1 points out a necessary condition involving real quadratic forms which Example 1 shows is not sufficient. Theorem 2 gives a new sufficient condition involving complex quadratic forms which Example 2 shows is not necessary.

*This work was done while the author was a National Academy of Sciences National Research Council Postdoctoral Research Associate at the National Bureau of Standards, Washington, D.C. 20234.

[†]Present Address: Institute for Fluid Dynamics and Applied Mathematics, University of Maryland, College Park, Maryland 20850.

THEOREM 1. Suppose $A \in DL_n(R)$. Then for each $0 \neq x \in R^n$, there is a $D \in D_n$ (which depends on x) such that

$$x^T D A x \geq 0.$$

Proof. Suppose to the contrary that there is a nonempty (necessarily open) set $S \subset R^n$ such that for all $D \in D_n$, $x \in S$

$$x^T D A x < 0. \quad (*)$$

Suppose notationally that

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \quad Ax = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}, \quad \text{and} \quad D = \begin{bmatrix} d_1 & & 0 \\ & \ddots & \\ 0 & & d_n \end{bmatrix}.$$

Since S is open, we may choose $x \in S$ such that $x_i y_i \neq 0$, $i = 1, \dots, n$. Our assumption $(*)$ now becomes

$$\sum_{i=1}^n y_i d_i x_i < 0 \quad (**)$$

for all choices $d_1, \dots, d_n > 0$. Thus it must be that each $y_i x_i < 0$, $i = 1, \dots, n$. It then follows that for any negative number λ , the system

$$d_i y_i = \lambda x_i, \quad i = 1, \dots, n,$$

may be solved by choice of $d_1, \dots, d_n > 0$. Such a solution is equivalent to the statement

$$DAx = \lambda x,$$

which means $\lambda \in \sigma(DA)$. The existence of a negative member of $\sigma(DA)$ contradicts $A \in DL_n(R)$ and completes the proof.

Note. In the statement of Theorem 1, " \geq " may not be replaced by " $>$ "

since $\begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix} \in DL_2(R)$.

Let $\overline{D_n}$ denote the Euclidean closure of D_n . By a straightforward compactness argument, it follows from Theorem 1 that

COROLLARY 1.1. *If $A \in \overline{DL_n(R)}$, then for each $0 \neq x \in R^n$ there is a $D \in \overline{D_n}$ (which depends on x) such that $x^T D x > 0$ and $x^T D A x \geq 0$.*

In a different context Fiedler and Ptak have studied the class defined by the necessary condition of Corollary 1.1 and related it to the class P_0 which is defined as follows [2].

DEFINITION. A is an element of the class P_0 if and only if all the principal minors of $A \in M_n(R)$ are nonnegative.

LEMMA. [2] *For $A \in M_n(R)$, the following are equivalent:*

- (i) $A \in P_0$;
- (ii) *for each $0 \neq x \in R^n$, there is a $D \in D_n$ (depending on x) such that $x^T D x > 0$ and $x^T D A x \geq 0$;*
- (iii) *each real eigenvalue of each principal submatrix of A is nonnegative.*

It then follows from Corollary 1.1 and the lemma that

COROLLARY 1.2. $\overline{DL_n(R)} \subseteq P_0$.

This containment is implicit in [7].

EXAMPLE 1. The converse of Theorem 1 is false (as well as those of the corollaries). Let

$$A = \begin{bmatrix} 1 & 0 & 27 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}.$$

Then $4 \in \sigma(A)$, and since $\text{Tr}(A) < 4$ $A \notin \overline{DL_3(R)}$. However, since all prin-

cipal minors of A are positive, it follows from Theorem 1.1 of [2] that for each $0 \neq x \in R^n$, there is a $D \in D_n$ such that

$$x^T D A x > 0.$$

Although the containment in Corollary 1.2 is proper, it suggests an alternative to studying $\overline{DL_n(R)}$ would be found in the following class.

DEFINITION. Let $P_0^- =$

$$\{A \in P_0: \text{there is a } \lambda \in \sigma(A) \text{ with } \operatorname{Re}(\lambda) < 0\}.$$

The lemma points out that if $A \in P_0$ and $\operatorname{Re}(\lambda) < 0$ for some $\lambda \in \sigma(A)$, then $\operatorname{Im}(\lambda) \neq 0$.

REMARK. $\overline{DL_n(R)} = P_0 - D_n P_0^-$.

Thus to characterize $\overline{DL_n(R)}$ it would suffice to obtain a satisfactory characterization of P_0^- .

THEOREM 2. If $A \in M_n(C)$ is such that for each $0 \neq x \in C^n$ there is a $D \in D_n$ (depending on x) such that

$$\operatorname{Re}(x^* D A x) > 0,$$

then $A \in DL_n(C)$.

Proof. Suppose $A \notin DL_n(C)$. Then there is an $E \in D_n$ such that there is a $\lambda \in \sigma(EA)$ with $\operatorname{Re}(\lambda) \leq 0$. Suppose

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

is such that

$$EAx = \lambda x,$$

and let

$$Ax = y = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}.$$

It then follows that $\operatorname{Re}(y_i \bar{x}_i) \leq 0, i = 1, \dots, n$. Now

$$x^* D A x = \sum_{i=1}^n d_i y_i \bar{x}_i,$$

where $D = \operatorname{diag}\{d_1, \dots, d_n\}$ and the right-hand side clearly has nonpositive real part for all $D \in D_n$. The statement of the theorem is the contrapositive of what has been shown.

By continuity we have also that

COROLLARY 2.1. *If $A \in M_n(C)$ is such that for each $x \in C^n$ there is a $D \in D_n$ (depending on x) such that*

$$\operatorname{Re}(x^* D A x) \geq 0,$$

then $A \in \overline{DL_n(C)}$.

EXAMPLE 2. The converses of Theorem 2 and Corollary 2.1 are false. Let

$$A = \begin{bmatrix} 1 & 0 & -150 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \quad \text{and} \quad x = \begin{bmatrix} -9 + 20i \\ -4 - 4i \\ 4 + 2i \end{bmatrix}.$$

Then it may be verified [3, 6] that $A \in DL_3(C)$. However, if

$$Ax = y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix},$$

then

$$\operatorname{Re}(y_i \bar{x}_i) \leq 0, \quad i = 1, 2, 3, \text{ and } \operatorname{Re}(y_i \bar{x}_i) < 0, \quad i = 1, 2.$$

Thus

$$\operatorname{Re}(x^* D A x) < 0$$

for all $D \in D_n$.

REFERENCES

1. K. J. Arrow and M. McManus, A note on dynamic stability, *Econometrica* **26**, 448–454 (1958).
2. M. Fiedler and V. Ptak, Some generalizations of positive definiteness and monotonicity, *Numerische Mathematik* **9**, 163–172 (1966).
3. D. Hook, unpublished communication, 1972.
4. C. R. Johnson, Matrices whose hermitian part is positive definite, thesis, California Institute of Technology, 1972.
5. C. R. Johnson, Gersgorin sets and the field of values, *J. Math. Anal. and Appl.* **45**, 416–419 (1974).
6. C. R. Johnson, Second, third, and fourth order D -stability, *J. Research NBS*, Section B, **78B**, 11–13 (1974).
7. J. Quirk and R. Ruppert, Qualitative economics and the stability of equilibrium, *Rev. Ec. Studies* **32**, 311–325 (1965).
8. J. Quirk and R. Saposnick, *Introduction to General Equilibrium Theory and Welfare Economics*, McGraw-Hill, New York (1968).
9. O. Taussky, A remark on a theorem of Lyapunov, *J. Math. Anal. and Appl.* **2**, 105–107 (1961).

Received May 16, 1973; revised August 29, 1973